2021 Michigan State Wheat Performance Trials

Photo: Fusarium Head Blight Nursery, 2021







2021 Michigan State Wheat Performance Trials

Dennis Pennington, Eric Olson, Sam Martin, Amanda Noble, Amelia Orr July 29, 2021

Cooperating weather last fall allowed growers to get wheat planted on time. Most of the crop across the state emerged, tillered out well and went into the winter in good shape. Temperatures over the winter were mild early, followed by cold temperatures in February. Nearly all of the wheat in the lower peninsula was protected by a blanket of snow. Little to no winterkill was reported. As the crop greened up in the spring, expectations and yield potential was very high. Spring conditions allowed nitrogen applications to be made timely, followed by fungicides at Feekes 9 and 10.5.1.

Dry conditions were welcomed for spring planting operations, however the dryness persisted resulting in 78% of the lower peninsula in D1 (moderate) or D2 (severe) drought. Many parts of the state received little or no rain during early to mid grain fill. Hardest hit areas and those with sandy soils caused the wheat to desiccate prematurely, ending grain fill. In late June, heavy and frequent rains made their way into the state. By the last week of June, the drought was over for much of MI. The Saginaw Valley and thumb regions were hardest hit by the drought. The premature death of the crop followed by heavy rains caused significant grain quality issues, particularly preharvest sprouting.

Falling number (FN) evaluates the degree of for preharvest sprouting. All white wheat production regions had poor FN to some degree with about 25% of the white wheat crop in MI below 240. Black sooty mold was widely reported across the state. Some areas had small amounts of black point. Conditions at flowering were dry, so vomitoxin levels were low in 2021. Persistent rains delayed harvest which worsened preharvest sprout. Even soft red wheat was at risk of sprout before the end of harvest.

Temperatures across the region were similar to '19. We did not have the excessive heat in '21 compared to '20. Rainfall was very short in April and May. Most of the rain in June came in two big rain events in the latter half of the month – near the end of the grain fill period.

Figure 1. Number of days above 90 F, 85 F and rainfall data from Michigan Automated Weather Station Network, MSU for three of the MSU Wheat Variety Trial Locations for the 2019, 2020 and 2021 growing seasons. 2021 data was reported through July 16, 2021.

		2019			2020			2021	
	Pigeon	Richville	Mason	Pigeon	Richville	Mason	Pigeon	Richville	Mason
Above 90 F	1	3	3	10	13	10	2	4	2
Above 85 F	14	12	16	30	33	30	15	16	19
April (in)	3.8	2.3	2.9	2.2	2.1	2.6	1.8	0.7	1.5
May (in)	2.8	5	3.4	3.3	3.8	4.2	1.2	1.2	2.6
June (in)	3.6	7	4.5	1.9	1.4	5.8	1.9	4.9	7.0
July (in)	1.9	2.4	2.3	2.8	3.2	2.1	2.5	1.0	1.5

Choosing Varieties

Variety selection is best made using at least three years of data. Varieties selected using data across all locations and multiple years will likely perform well under a wide range of conditions; although, performance of a given variety will vary based on testing location. In selecting varieties for a specific location, it is important to identify varieties that perform well near the location where the variety will be grown. Table 1 provides information on which varieties are top performers in each of the seven trial locations in 2019 through 2021. Selection and planting of two or more varieties is recommended. As an example, planting varieties that differ in flowering date can allow for staggering of management applications, specifically, fungicides to control Fusarium head blight. When selecting varieties, look at disease resistance as well as yield potential.

Disclaimer: MSU makes no endorsement of any wheat variety or brand.

Experimental Design

The 2021 State Wheat Performance Trial entries were planted in 7 counties: Gratiot, Hillman, Ingham, Huron, Lenawee, Sanilac and Tuscola. Appendix A (below) presents information on each of these sites. Each plot contained 6 rows with 7.5" row spacing and was planted to a length of 18 feet. Plots were trimmed to a length of 12 feet long in the spring for harvesting purposes. Sites were designed as Alpha Lattice with three replications. All seed was treated, but the chemicals and rates used varied according to the preferences of the originating organization. Seeding rates per linear foot of row were standardized to the rate that would equate with a stand of 1.8 million seeds per acre in a solid stand planted in 7.5" rows. Fall fertilizer application varied with cooperator practice. Spring nitrogen was applied as urea (90 lbs/acre actual N) at green-up and Affinity BroadSpec was used for weed control at all sites.

All sites were coordinated under high management with the exception of additional conventionally managed trials at Tuscola and Isabella Counties. Under high management, an additional 30 pounds of nitrogen was applied using streamer nozzles and 28% UAN. Quilt Xcel fungicide was tank mixed with herbicide and applied at Feekes 6. Prosaro fungicide was applied to control late season fungal diseases with application coinciding with the average flowering date of the trial location.

All plots within a location were harvested on a single day. Yield was calculated using the entire area of the plot including the wheel tracks between plots leading to an underestimation of yield. For data reported on a 0-9 scale 0 is the best possible score.

Seven of our experimental sites are on private farmland. We are extremely grateful to those growers for accommodating our work and all of the associated inconveniences. Funding for the high-management trial inputs was provided by the Michigan Wheat Program. Questions and comments regarding the research reported here should be directed to Dennis Pennington at pennin34@msu.edu or (269) 832-0497. This report and previous reports, may also be accessed through the Web at http://www.varietytrials.msu.edu/wheat.

Multi-Year Performance Summary

The full trial included 114 entries (52 of which were experimental lines) from 16 organizations, including Michigan State University, and data analyses were conducted using <u>all</u> of these entries. Attached to this narrative is a list of the names and contact information for those organizations. Each row in these tables has data for a single entry. The columns contain averages for a given trait and time period. Data for all of the entries in this trial are not presented here. However, the averages and statistical parameters in this report are based on the entire set of evaluated materials. <u>Comparisons among entries are only valid within a column</u>. Tables 1 and 2 are sorted first by grain color, and then in descending order by yield for 2021. Tables 3, 4 and 5 are sorted in alphabetic order by company and entry name. In some instances (e.g. yield), data columns to the right of the 2021 data columns are multi-year averages. Only data for entries included in all of the relevant years' tests are found here. Not all entries have been tested in all years, so the tables have several blank cells. See the section titled 'Experimental Design' for details on how the trials were conducted and for more detail on what the data in each column represents.

At the bottom of most columns in the tables is the trial average (mean), LSD (least significant difference), and CV (coefficient of variation) for data in that column. LSD values vary among traits and data sets (combinations of sites and years). Differences between the means for two entries that are greater than the LSD for that column are very likely to reflect a genuine difference between the two varieties. If the difference between two means is smaller than the LSD for that column, one should conclude that there is **no evidence that those entries are different for that trait** in the years and sites considered.

<u>Table 1</u> contains yield data. This data was acquired electronically on the plot combine at the time of harvest. Yield data is standardized to 13.5% moisture. The 2021 yield data contains the multi-site yield averages of only the high management sites and does not include the conventionally managed yield data from Tuscola and Isabella Counties. The conventionally managed data can be found in Table 4 in the conventional vs. high management results.

<u>Table 2</u> contains test weight and percent moisture for all locations along with the overall average across locations.

<u>Table 3</u> contains data on resistance to Fusarium Head Blight (FHB, scab). The 2020 deoxynivalenol (DON, VOM) numbers are reported. Once 2021 data from the lab are back, this report will be updated. Scab data were obtained from heavy disease pressure in an inoculated scab screening nursery. FHB infected grain is spread to provide inoculum and artificial misting provides disease-promoting conditions throughout the entire flowering period. 2021 grain samples will be submitted for DON analysis and will be reported later. **Cephalosporium stripe** ratings were taken at the Ingham County location. **Preharvest sprouting (PHS)** samples were collected from Ingham County and subjected to misting in the greenhouse for three days and rated for the degree of sprouting. PHS ratings were conducted using at 0-9 scale with 0 having no sprouting and 9 having fully emerged radicle and roots from over 80% of the spike.

The **flowering date** indicates the average number of days past January 1st that a given entry reached the point where ½ of its heads were flowering. **Physiological maturity** was recorded as the date when 50% of the peduncles in a plot were turning yellow. **Plant height** is reported as the distance in inches from the ground to the tip of average heads in a plot.

FHB Resistance Traits

Severity: The average percent of infected spikelets in each head. Incidence: The percent of all spikes in a plot showing infection. FHB index: The overall infection considering severity and incidence.

DON: Levels of mycotoxin (ppm) present in grain. DON data is from the 2020 crop year.

Levels of DON and severity are the most reliable traits to be used in selecting FHB-resistant varieties.

High Management vs. Conventional Management Performance

<u>Table 4</u> provides a comparison of variety performance under intensive management and conventional management practices. Data on yield, test weight, grain moisture at harvest are provided from conventional management and high management trials at Tuscola and Isabella Counties. Conventional management received 90 pounds of N per acre only. The high management received an additional 30 pounds of N per acre applied at Feekes 6 plus Quilt Xcel fungicide at Feekes 6.0, followed by Prosaro fungicide applied at Feekes 10.5.1. The last two columns presents the yield advantage of high management in bushels per acre as well as a ranking of the response. A positive number indicates a yield response to high management. A negative number indicates the higher management actually produced a lower yield. Overall means were 6.7 and 8.1 bushels per acre higher for the high management treatment at Tuscola and Isabella respectively.

Milling and Baking Quality

Table 5 contains data for milling and baking quality. Quality data are from the 2020 harvest season and prior. Data were generated by the USDA Eastern Soft Wheat Quality Laboratory in Wooster, Ohio on grain harvested from the Michigan State Variety trial each year. Flour yield is the ratio of the weight of extractable flour to the weight of milled grain, expressed as a percentage. Percent protein in flour is adjusted at 14% moisture. Softness equivalent percent is the softness of the flour, with higher values indicating softer grained wheat. For cookie diameter, a larger diameter is better. Whole grain protein (%) and whole grain hardness are being reported with 0-100, and higher values indicating harder wheat. The quality lab test weight is not identical to the test weight at harvest due to grain drying and grain cleaning prior to quality laboratory test weight evaluation. Solvent Retention Capacity (SRC) can be conducted on flour using several different solvents and reflects different characteristics of flour quality. Soft wheat flour for cookies typically have a target of 95% or less when used by the US baking industry for biscuits and crackers. Sodium carbonate SRC increases as starch damage due to milling increases. Normal values for good milling soft varieties are 68% or less. Lactic acid measures gluten strength with "weak" soft varieties having values below 85% and strong gluten soft varieties having values, typically, above 105% or 110%.

Table 1: Multi-Year Performance Summary (Note: Tables sorted by 2021 High Management Yield, white wheat's grouped before red)

		•	i i				ivialiagelli		_		I		naham		ı —		Human		ı —		Camilaa			т.	.aaala		Lana		Jackelle
Lina	Seed	Awns	Chaff		u/A adjus	sted to 13.5%			<u>Hillma</u>		۱	_	ngham	24.4	۱		<u>Huron</u>	24.4	۱ .,		Sanilac 2 V A	27.4			uscola 2 V		<u>Lena</u>		<u>Isabella</u>
Line	Color	AWIIS	Color	2021		2 Yr Avg	3 Yr Avg	20		2 Yr Avg)21	2 Yr Avg	3 Yr Avg	20		2 Yr Avg	3 Yr Avg		021	2 Yr Avg	3 Yr Avg	20		2 Yr Avg	3 Yr Avg	202		2021
				Overall	Rank	20-21	19-21	Bu/A	Rank	20-21	Bu/A	Rank	20-21	19-21	Bu/A	Rank	20-21	19-21	Bu/A	Rank	20-21	19-21	Bu/A	Rank	20-21	19-21	_	Rank	Bu/A Rank
DF 271 W	W	Awnletted	W	97.4	1			91.9	18		81.0	1			80.1	3			101.4	8			111.4	2			107.1	3	108.1 2
AgriMAXX Exp. 2050W	W	Awnletted	W	97.2	2	97.8		96.5	7	104.5	71.4	16	81.3		74.7	10	92.6		107.6	1	104.1		114.7	1	106.5		106.9	4	109.2 1
Jupiter	W	Awnletted	В	94.8	3	93.4	97.0	87.9	28	97.0	73.9	10	79.5	82.5	82.1	1	95.7	101.0	103.0	3	97.0	100.3	100.5	7	98.0	104.1	109.8	2	106.7 4
KWS317	W	Awnletted	W	94.7	4	99.8		97.4	3	108.7	76.7	3	88.9		75.1	9	93.1		102.8	4	108.8		99.0	8	99.2		106.5	6	104.4 7
ISF 1115	W	Awnletted	W	94.0	5			93.7	13		76.3	6			76.0	7			96.7	17			102.0	4			103.9	8	107.5 3
MI17W0235	W	Awnletted	В	92.0	6	91.4		90.8	21	95.1	76.5	4	84.6		69.2	20	86.7		96.6	18	91.9		100.9	6	98.6		103.8	9	106.6 5
MI17W0224	W	Awnletted	W	91.4	7	94.2		86.3	29	99.1	72.5	13	76.1		79.5	4	96.4		99.1	10	98.2		102.1	3	101.2		101.3	14	98.2 23
MI16W0133	W	Awned	W	91.3	8	92.8	96.4	90.5	22	100.4	74.6	9	83.4	87.0	73.5	11	88.9	97.7	95.2	22	96.2	98.6	92.5	21	95.0	102.3	111.1	1	103.3 10
Whitetail	W	Awnletted	W	90.9	9			88.8	26		73.3	12			77.4	5			90.5	29			96.9	14			102.5	10	104.0 9
MI14W0190	W	Awnletted	В	90.7	10	89.9	91.9	94.1	11	95.6	76.2	7	81.6	84.6	70.4	18	86.3	92.0	102.4	7	94.6	94.0	90.3	24	91.2	96.9	99.5	19	100.5 17
MI16W0528	W	Awnletted	W	90.7	11	92.7	95.7	90.3	23	99.6	76.3	5	82.6	84.7	70.4	17	88.5	96.3	97.0	16	95.9	97.6	97.7	11	96.7	104.2	102.2	12	100.1 20
ISF 1001	W	Awnletted	W	90.2	12			91.5	19		71.5	15			76.8	6			102.8	5			93.4	20			96.0	26	100.2 19
AgriMAXX EXP 2055W	W	Awnletted	W	90.1	13	91.9		97.2	6	102.3	71.8	14	78.5		67.6	25	86.4		99.3	9	96.1		95.3	17	96.0		101.9	13	97.1 25
DF 261 W	W	Awnletted	W	90.0	14			94.4	10		70.4	18			68.4	21			102.7	6			97.5	12			98.2	21	98.7 22
KWS308	W	Awnletted	W	89.9	15			97.7	2		75.8	8			61.0	29			97.3	14			101.4	5			98.0	22	97.5 24
MI18W0286	w	Awnletted	w	89.8	16			89.6	25		70.5	17			72.4	13			95.3	21			98.3	9			102.4	11	100.4 18
Dyna-Gro WX21791W	w	Awnletted	w	89.8	17			95.2	9		65.9	27			72.3	14			94.2	25			94.7	19			100.6	16	104.1 8
KWS319	w	Awnletted	w	89.7	18			92.6	15		68.1	25			67.7	24			103.3	2			96.7	15			96.4	25	101.3 13
Ambassador	w	Awnletted	w	89.4	19	91.2	95.3	97.4	4	101.4	73.6	11	76.1	81.4	75.5	8	91.9	100.5	94.9	24	96.0	101.7	87.1	28	90.7	97.9	97.4	23	100.5 16
MI18W0200	w	Awnletted	w	89.3	20			97.3	5		68.2	24			72.1	16			98.1	11			95.0	18			92.3	29	101.9 12
DF 218 W	w	Awnletted	w	89.2	21	92.7	93.9	95.5	8	105.4	76.9	2	84.1	86.2	65.0	26	87.0	94.7	93.9	26	91.0	92.6	95.7	16	96.1	102.3	96.7	24	100.9 15
Moonlight	w	Awnletted	w	88.8	22	90.0	92.9	92.4	16	98.3	70.3	20	73.9	79.4	72.3	15	89.9	96.5	93.8	27	90.3	92.3	97.0	13	97.4	103.3	100.8	15	98.8 21
KWS327	w	Awnletted	w	88.4	23			99.6	1		69.9	22	75.5		62.9	28			97.2	15			97.7	10			98.4	20	92.3 29
Dyna-Gro 9242W	w	Awnletted	w	88.4	24	91.0	94.5	90.3	24	103.4	68.7	23	77.7	84.8	69.6	19	87.4	95.5	95.7	20	92.4	94.6	90.2	26	94.2	103.0	99.9	17	103.2 11
Dyna-Gro WX19799W	w	Awned	w	88.3	25	J1.0	J4.J	88.1	27		70.3	19			67.8	23			93.7	28	J2.4	J4.0 	90.3	25	J4.Z		106.4	7	100.9 14
Dyna-Gro 9082W	w	Awned	w	88.2	26	86.4		93.9	12	100.0	62.5	29	66.9		63.0	27	80.9		96.0	19	99.1		89.9	27	84.8		106.4	5	106.1 6
KWS305	w	Awnless	w	87.9	27			93.5	14		64.2	28			80.2	21			97.3	13			91.6	23			93.9	28	96.8 26
KWS316	w	Awnletted	w	87.9	28	89.9		92.3	17	97.6	70.0	21	76.8		73.1	12	89.5		94.9	23	91.9		92.3	22	93.6		95.8	27	95.8 28
AC Mountain	W	Awnletted	w	86.8	29	86.3	88.7	91.0	20	94.0	66.9	26	74.6	76.9	68.3	22	86.3	92.2	97.9	12	86.6	87.9	86.8	29	89.8	98.0	99.7	18	96.5 27
ISF 790	R	Awnletted	W	96.7	1			97.2	6		74.9	22	74.0		86.4	1			99.3	24			104.6	0			104.6	36	109.7 4
	R	Awned	w		2				-		77.9	22			84.7				100.3	16			107.3	2			105.0		
DF 121 R	R		W	96.5		02.0		88.8	55	00.0		3	00.2			2	00.1			16	07.6			3				30	110.4 2
LW2068	R	Awned	W	94.8 94.7	3	92.8		89.3 97.3	53 5	96.9	77.3	5	80.3		75.2	17	90.1		102.6 105.1	3	97.6		102.7	16	99.1 104.7		108.0	11	107.7 16 108.2 11
MCIA MARLIN Haubert	R	Awnletted	W	93.9	-	97.2		97.8	4	111.4	71.8 77.1	48 7	76.0		78.0 71.8	5 37	98.7		103.1	5	95.2		96.3 98.6	47 36	104.7		105.3 102.7	28 48	108.2 11 104.4 41
MCIA Wharf	R	Awned Awnletted	W	93.9	5	96.8	100.4	95.6	12	103.7	77.8	/	85.1	88.3	71.8	37	92.0	99.4	95.6	46	101.8	103.2	105.3	7	101.5	110.5	102.7	48	104.4 41
					-		100.4					4		00.3		39		99.4		_		105.2				110.5			
DF 131 R	K	Awned	W	93.8	7	93.9		91.1	42	98.7	75.0	21	82.6		77.3	7	93.2		94.4	51	99.2		101.2	23	99.7		111.1	2	106.2 25
W 322	R	Awned	W	93.7	8			94.3	22		71.4	50			74.7	24			97.6	38			102.8	14			110.8	3	107.9 12 107.8 14
RWEXP1212	R	Awnletted		93.7	9			96.9	7		72.8	36			77.2	8			95.6	44			100.2	28			105.0	30	
REXP1215	R	Awned	W	93.5	10			85.4	66		78.5	2			66.2	60			104.0	_			108.0	1			106.8	17	103.9 48
AgriMAXX 498	R	Awnletted	W	93.4	11	93.5		90.2	48	101.2	79.7	1	85.7		75.1	18	88.5		106.5	2	93.6		96.2	49	98.4		101.8	56	105.3 33
MI16R0906	R	Awnletted	W	93.4	12	96.7	07.6	88.7	57	99.4	76.8	10	83.4		76.1	13	93.0	101.7	97.2	41	106.9	100.0	103.6	11	100.8	105.0	106.4	20	104.8 37
MCIA Flipper	K	Awnletted	W	93.3	13	96.1	97.6	94.5	21	108.2	72.1	43	78.8	81.7	66.2	60	90.9	101.7	107.8		101.7	100.9	99.6	33	100.6	105.9	104.0	38	108.6 7
Dyna-Gro 9002	R	Awned	W	93.2	14	93.9	96.5	95.5	13	103.0	75.1	20	80.4	83.6	72.1	33	91.1	99.3	100.3		98.6	98.9	96.5	44	96.3	104.2	109.7	7	105.8 27
Dyna-Gro 9172	R	Awned	W	93.1	15	94.2		98.0	2	103.8	72.3	41	82.2		65.3	66	88.7		98.7	30	97.2		102.4	17	99.0		111.3	1	107.2 20
Tyson	R	Awned	W	93.1	15			94.8	16		72.4	40			71.2	41			97.7	36			100.0	29			108.2	9	108.6 9
W 324	R	Awned	W	93.1	17			93.2	30		76.8	11			63.8	69			93.4	56			102.7	15			107.9	12	112.4 1
Dyna-Gro 9070	R	Awned	W	93.1	18	95.6	98.2	84.0	69	97.0	74.8	26	78.3	83.1	74.8	22	92.9	99.4	98.9	27	106.3	101.9	106.9	4	103.3	108.5	105.2	29	107.6 17
RS 912	R	Awnless	W	93.0	19	95.9		91.9	39	104.9	73.1	35	81.8		72.5	31	90.7		100.0		101.4		104.1	9	100.5		106.9	15	103.6 54
MI17R0357	R	Awnletted	W	92.9	20	98.2		93.4	29	105.7	74.1	27	74.9		75.9	15	97.7		92.8	61	109.1		103.1	13	103.5		103.2	45	106.7 21
DF EX 2103 R	R	Awned	W	92.9	21			98.2	1		72.5	39			64.5	67			99.9	20			97.5	39			108.5	8	110.3 3
KWS340	R	Awnletted	W	92.9	21			98.0	2		76.3	13			75.3	16			99.1	25			93.2	56			102.6	49	106.1 26
KWS361	R	Awnless	W	92.8	23			90.1	49		71.6	49			76.4	10			95.3	47			107.7	2			102.1	54	105.3 31
Dyna-Gro WX20734	R	Awned	W	92.8	24			86.3	65		76.2	16			76.2	12			98.8	29			100.7	25			107.3	14	105.4 29
SY 100	R	Awnletted	W	92.7	25	95.1	100.0	94.6	19	102.3	75.5	17	81.1	88.4	68.1	54	87.0	98.3	94.2	53	102.8	102.9	106.0	5	102.3	110.3	103.6	42	105.0 36
RS 902	R	Awned	W	92.6	26	92.4	96.8	90.3	46	102.7	71.8	47	81.6	85.9	71.0	44	89.3	99.2	103.2	6	92.7	96.5	96.4	45	95.7	105.6	110.2	5	107.4 19
AgriMAXX 513	R	Awned	W	92.6	27			92.2	37		76.3	14			74.1	27			99.0	26			97.3	41			105.7	23	104.6 39
AgriMAXX 505	R	Awned	W	92.6	28	94.1		91.1	43	101.2	72.1	45	80.7		71.9	35	90.1		99.3	23	100.4		101.9	18	98.1		106.9	16	104.2 44
W 304	R	Awned	W	92.5	29	93.1	95.6	95.9	11	103.8	77.2	6	84.5	87.4	72.1	34	90.7	99.9	95.6	45	92.1	91.9	93.4	55	94.4	103.0	105.5	26	106.3 23

Table 1: Multi-Year Performance Summary (Note: Tables sorted by 2021 High Management Yield, white wheat's grouped before red)

Table 1 : Multi-Year Per	Torman	ce Summary	(Note: I	ables sor	rtea by	ZUZI HIgh	ivianagem	ient Yie	_		s grou	oea bei	rore rea)															
	Seed		Chaff	Yield (B	Bu/A adjus	ted to 13.5%			Hillm	<u>an</u>		<u>lı</u>	ngham				<u>Huron</u>			9	<u>Sanilac</u>			<u>T</u>	<u>'uscola</u>		<u>Lenawee</u>	<u>Isabella</u>
Line	Color	Awns	Color	2021		2 Yr Avg	3 Yr Avg	20	21	2 Yr Avg	20	21	2 Yr Avg	3 Yr Avg	20	21	2 Yr Avg	3 Yr Avg	202	21	2 Yr Avg	3 Yr Avg	202	1	2 Yr Avg	3 Yr Avg	2021	2021
	COIO		COIO	Overall	Rank	20-21	19-21	Bu/A	Rank	20-21	Bu/A	Rank	20-21	19-21	Bu/A	Rank	20-21	19-21	Bu/A	Rank	20-21	19-21	Bu/A	Rank	20-21	19-21	Bu/A Rank	Bu/A Rank
RS 977	R	Awned	W	92.4	30	93.3	96.1	92.5	32	101.0	76.6	12	86.8	88.3	70.5	46	86.9	96.5	102.0	10	99.4	96.4	94.6	52	92.6	103.4	107.5 13	103.8 49
LW2169	R	Awned	W	92.2	31			90.6	44		68.7	64			68.0	55			101.4	11			98.6	35			110.3 4	108.6 8
LW2148	R	Awnletted	W	92.2	32			95.1	14		73.7	29			72.5	32			96.0	43			99.9	31			103.6 43	103.0 60
AgriMAXX 516	R	Awned	W	92.1	33			94.7	17		68.9	61			73.3	30			95.0	48			100.0	30			105.0 32	107.8 13
HS338R	R	Awnletted	W	92.1	34	93.7		92.4	34	103.1	76.8	9	81.5		69.1	52	88.0		101.1	12	96.2		97.2	42	99.8		102.5 51	104.7 38
Dyna-Gro 9182	R	Awnless	W	92.0	35	93.3		94.2	23	104.8	72.8	37	81.5		74.6	25	90.9		99.6	21	95.9		94.5	53	93.7		102.1 53	104.5 40
Sunburst	R	Awnless	W	92.0	36	92.3		92.3	36	102.0	70.7	55	82.5		75.0	20	88.9		99.6	22	92.2		96.3	46	95.7		105.7 24	103.5 56
MI16R0720	R	Awnletted	W	91.9	37	91.9		94.9	15	102.0	71.1	52	73.1		71.5	40	92.0		94.8	50	89.0		103.4	12	103.2		106.5 19	101.0 64
RS 961	R	Awnletted	W	91.8	38	94.0	97.1	93.6	27	99.2	74.9	25	83.9	90.2	76.0	14	89.8	96.5	100.5	13	107.4	103.6	85.0	71	89.9	98.2	101.7 57	109.5 5
Dyna-Gro 9151	R	Awned	W	91.8	39	93.9		88.9	54	101.0	76.2	15	84.6		65.8	63	85.9		98.9	28	101.3		100.6	26	96.6		108.1 10	105.4 29
W 310	R	Awned	W	91.7	40	93.6		87.4	62	98.9	73.7	30	86.7		69.9	50	86.8		96.6	42	95.8		103.8	10	99.8		106.8 18	102.3 62
AgriMAXX 502	R	Awned	W	91.7	41			88.8	55		70.9	54			77.1	9			97.3	40			101.4	22			101.5 59	103.4 58
MI16R0898	R	Awnletted	W	91.5	42	91.8	96.4	83.6	70	92.4	75.3	19	80.8	84.9	74.0	28	87.8	94.6	102.0	9	103.6	105.6	93.0	57	94.6	100.4	106.2 22	103.8 49
DF EX 2104 R	R	Awnletted	W	91.4	43			93.5	28		69.6	57			80.7	3			98.6	32			99.0	34			95.1 69	103.6 52
SY Viper	R	Awnletted	W	91.2	44	95.5	96.5	87.1	63	103.7	73.4	33	80.6	81.3	77.8	6	95.1	102.1	97.8	35	101.9	99.5	90.5	65	96.3	103.2	103.2 46	106.4 22
LW2958	R	Awned	W	90.9	45			92.4	35		72.6	38			73.6	29			94.4	52			95.1	51			103.7 41	103.0 59
W 305	R	Awnletted	W	90.9	46	94.6	98.0	89.6	50	96.5	77.0	8	86.2	90.3	67.7	57	86.0	94.7	90.4	65	104.1	101.2	101.8	19	100.4	105.6	105.5 25	105.1 35
MCIA Jonah	R	Awnletted	W	90.9	47	92.4	96.6	89.3	52	101.4	73.6	31	82.9	87.1	74.5	26	88.8	99.1	97.7	37	92.2	96.2	91.7	60	96.7	103.9	105.3 27	103.4 57
MI18R1605	R	Awnletted	W	90.9	48			96.1	9		70.6	56			74.8	22			92.6	62			99.7	32			99.9 62	104.0 47
DF 112 R	R	Awned	W	90.7	49	89.8	93.8	92.1	38	102.2	69.3	59	75.5	80.6	78.9	4	91.8	100.8	87.3	69	82.6	88.4	97.0	43	97.0	105.3	106.3 21	105.5 28
DF EX 2101 R	R	Awned	W	90.6	50			88.0	59		64.7	71			70.4	47			97.3	39			100.4	27			109.8 6	105.3 33
DF 105 R	R	Awned	W	90.4	51	92.0	94.1	90.5	45	102.5	74.9	22	82.8	86.8	74.9	21	94.3	97.0	87.8	67	83.0	88.3	97.5	38	97.4	104.3	104.9 34	101.7 63
SY 547	R	Awnless	W	90.3	52	94.6	95.7	93.9	25	104.9	68.7	62	78.6	83.2	65.9	62	86.3	92.9	91.5	63	105.4	102.4	101.6	20	98.1	104.4	102.6 50	108.4 10
MCIA Red Dragon	R	Awnless	W	90.1	53	89.5	90.8	94.1	24	105.7	73.8	28	75.6	76.5	71.1	43	87.5	97.5	85.8	71	81.8	83.9	101.1	24	96.8	105.0	96.3 66	107.7 15
LCS3334	R	Awnletted	W	90.0	54	89.9	92.1	87.9	60	94.0	74.9	22	82.3	82.7	75.1	19	90.2	97.2	98.6	32	87.6	87.5	96.2	48	95.1	100.9	93.7 70	104.0 46
AgriMAXX 503	R	Awnless	W	90.0	55	92.3		79.5	71	98.3	67.7	66	78.5		67.1	58	85.9		98.2	34	97.2		105.7	6	101.5		104.5 37	107.6 17
DF 119 R	R	Awnletted	W	90.0	56	94.5	99.3	94.7	17	102.3	69.0	60	74.0	82.4	70.0	48	92.8	102.1	91.5	63	102.3	105.5	97.9	37	101.0	107.0	100.9 61	104.0 45
AgriMAXX EXP 2002	R	Awned	W	90.0	57			93.8	26		71.0	53			71.9	36			94.0	55			95.1	50			101.2 60	103.6 51
W 313	R	Awnless	W	89.9	58	92.4		90.2	47	103.0	73.2	34	79.7		59.2	71	84.0		100.2	17	97.3		101.6	21	97.9		103.7 39	100.8 67
DF EX 2102 R	K D	Awned	W	89.9	59			89.4 96.0	51		73.5 75.3	32			68.4	53			93.1 93.2	59			97.3 92.8	40			101.9 55 99.5 63	105.3 32
Dyna-Gro WX20738 Dyna-Gro 9120	R D	Awned	W	89.1 89.1	60			87.4	10 61		69.4	18			64.2 65.7	68 64			100.1	58			91.5	58 61			99.5 63 103.7 40	103.6 53 106.3 24
MCIA Whale	P	Awned Awnletted	W	88.4	61 62	91.8	94.5	85.4	67	99.1	67.4	58 69	81.0	86.0	70.8	45	89.3	96.0	100.1	18 14	99.1	98.2	86.7	70	90.7	98.1	103.7 40	106.5 24
KWS356	R	Awned	W	88.4	63	91.0	94.5	88.1	58	99.1	71.4	51	61.0		66.7	59	09.5	90.0	94.1	54	99.1	90.2	92.6	59	90.7	96.1	104.6 35	104.4 42
Prestyn	R	Awned	W	88.4	63			93.0	31		72.2	42			71.2	42			93.3	57			90.7	64			95.4 68	100.9 65
DF 141 R	R	Awned	W	88.3	65	89.7		86.3	64	98.3	67.5	68	76.5		76.3	11	93.5		90.0	66	88.2		90.2	66	91.9		102.9 47	100.9 63
W 300	R	Awned	w	88.2	66			91.1	41		72.1	44			65.6	65			102.8	7			94.3	54			97.1 64	92.6 69
SY 576	R	Awned	W	87.7	67	87.5	91.3	91.8	40	97.5	68.7	63	75.8	80.1	67.9	56	84.4	91.3	93.0	60	93.0	94.5	91.0	63	86.9	99.2	101.6 58	100.3 68
KWS375	R	Awned	w	86.7	68			96.2	8		68.7	65			62.6	70			86.3	70			87.3	69			105.0 33	104.3 43
Harbor	R	Awnletted	w	86.5	69	88.4	90.9	92.5	33	101.9	67.5	67	72.6	77.7	69.9	49	87.5	96.1	98.7	31	87.4	90.6	87.5	68	92.7	99.3	97.0 65	91.9 70
13VTK59-55	R	Awnletted	w	86.3	70			84.7	68		66.7	70			71.8	38			87.6	68			91.4	62			96.1 67	103.6 54
Erisman	R	Awnletted	W	86.0	71	88.5		94.5	20	100.4	71.8	46	77.3		69.8	51	88.0		94.9	49	84.7		88.0	67	92.3		93.1 71	91.0 71
			Mean	91.2		92.8	95.3	92.1		101.1	72.6		80.1	84.0	71.9		89.7	97.3	97.3		96.4	96.7	97.6		96.9	103.2	103.2	103.8
			cv	2.6		3.0	2.9	4.2		3.7	4.9		5.3	4.4	7.1		3.9	3.1	4.6		6.9	6.0	5.9		4.5	3.4	4.3	4.0
			LSD	2.5		3.0	2.7	8.3		5.6	3.5		4.6	3.9	11.5		4.3	3.9	10.0		11.2	8.3	9.3		5.5	3.9	3.0	2.3
																			•									•

Table 2. Multi-Location Performance Summary for Test Weight and Percent Moisture.

Table 2. Multi-Location		Ove		Hilln		Ingh		Hur	on	San	ilac	Tusc	ola	Lenav	wee	Isabe	ella
Line	Color	% Moist	TW														
13VTK59-55	R	14.2	59.4	14.3	58.7	13.8	58.9	12.8	60.3	15.8	58.1	13.7	61.3	14.1	61.0	14.9	57.3
AC Mountain	W	14.0	57.0	14.4	58.2	12.8	55.8	13.2	58.0	15.9	56.3	13.3	57.8	13.6	58.0	14.8	55.0
AgriMAXX 498	R	14.9	57.5	16.8	57.9	14.7	57.5	13.1	58.3	16.0	56.9	14.4	59.0	14.2	57.7	14.9	55.4
AgriMAXX 502	R	14.3	57.9	14.3	58.4	13.9	56.2	13.2	59.9	15.8	56.7	13.9	59.6	14.0	59.1	15.0	55.4
AgriMAXX 503	R	13.9	57.3	11.1	53.6	14.7	55.7	13.5	59.4	15.3	57.0	14.2	60.2	13.9	59.2	14.6	56.4
AgriMAXX 505	R	14.1	59.3	14.6	58.7	14.3	58.0	12.5	61.2	15.4	58.3	13.6	61.4	14.0	61.0	14.6	57.1
AgriMAXX 513	R	14.2	58.5	15.2	58.2	14.5	56.7	12.3	60.3	15.6	57.4	13.7	60.7	13.7	59.6	14.5	56.7
AgriMAXX 516	R	14.4	57.7	15.1	58.6	14.0	55.3	13.4	59.8	15.5	57.0	14.1	59.3	14.3	58.4	14.3	55.8
AgriMAXX EXP 2002	R	14.4	59.0	15.1	58.9	14.5	57.3	12.9	61.1	14.9	58.2	14.0	60.8	14.0	59.8	15.1	57.5
AgriMAXX EXP 2055W	W	14.6	57.0	15.2	58.1	15.1	55.8	13.3	57.1	15.6	55.4	14.0	58.0	14.0	58.4	14.9	55.9
AgriMAXX Exp. 2050W	W	14.5	57.2	14.4	58.4	15.2	55.6	12.9	58.1	17.3	56.3	13.6	58.3	13.8	58.3	14.3	55.5
Ambassador	W	14.2	56.5	14.2	58.2	14.9	53.5	13.3	59.2	15.6	56.0	13.5	57.0	13.6	56.7	14.3	54.9
DF 105 R	R	14.1	57.6	14.1	58.1	14.7	55.9	12.7	60.1	15.0	57.2	13.7	58.8	13.9	57.5	14.7	55.6
DF 112 R	R	14.1	57.2	14.2	58.3	14.2	55.0	13.2	59.2	15.1	56.9	14.0	58.7	13.6	57.4	14.3	55.2
DF 119 R	R	14.4	57.9	14.7	58.5	14.8	55.9	12.7	60.0	15.6	56.1	13.9	59.6	13.7	58.2	15.1	56.7
DF 121 R	R	14.6	57.8	15.1	58.5	15.0	56.1	13.4	60.4	16.2	57.3	13.0	56.3	14.2	59.0	15.5	56.7
DF 131 R	R	14.4	58.0	15.5	58.7	14.4	55.7	12.9	60.0	15.8	56.9	13.7	59.7	13.9	58.3	14.6	56.8
DF 141 R	R	14.1	58.1	15.1	58.6	14.7	54.9	13.2	60.7	14.5	57.8	13.2	60.0	14.1	58.8	14.5	55.8
DF 218 W	W	14.8	58.0	17.0	58.0	15.1	56.3	12.6	60.4	15.7	57.7	14.0	60.0	14.1	58.1	15.2	55.7
DF 261 W	W	14.3	57.1	14.9	58.3	14.2	56.1	12.2	57.8	15.7	55.3	13.9	58.2	14.0	58.6	15.1	55.5
DF 271 W	W	14.2	57.4	14.5	57.9	14.7	56.8	13.2	58.2	15.6	55.5	13.4	59.4	13.7	59.3	14.3	55.0
DF EX 2101 R	R	14.1	58.1	15.3	58.4	13.7	56.4	12.4	59.5	14.9	56.7	13.5	60.7	13.5	59.7	14.8	55.7
DF EX 2102 R	R	14.1	58.3	14.6	58.0	14.6	56.4	12.5	60.4	13.8	57.2	13.7	60.1	13.9	59.2	15.2	57.2
DF EX 2103 R	R	14.2	56.2	15.2	57.9	13.4	54.6	12.2	57.3	15.8	55.4	14.0	57.1	14.0	57.0	14.3	53.9
DF EX 2104 R	R	14.3	59.1	14.7	59.0	14.5	56.3	12.9	61.1	15.8	58.2	13.8	61.0	14.2	60.2	14.6	58.3
Dyna-Gro 9002	R	14.5	57.8	15.4	58.3	14.1	56.8	12.9	59.4	16.3	57.2	14.2	59.1	13.9	58.8	14.5	55.6
Dyna-Gro 9070	R	14.2	57.9	14.4	58.4	14.6	56.6	12.8	58.8	14.7	56.6	13.8	60.1	13.8	58.8	15.1	55.6

Table 2. Multi-Location Performance Summary for Test Weight and Percent Moisture.

	renomian	Ove		Hilln		Ingh		Hur	on	San	ilac	Tusc	ola	Lenav	wee	Isab	ella
Line	Color	% Moist	TW														
Dyna-Gro 9082W	W	14.3	57.5	15.6	58.8	14.2	52.8	12.4	59.2	15.5	57.4	13.6	58.7	13.6	58.9	14.7	56.6
Dyna-Gro 9120	R	14.6	59.3	15.2	58.6	14.2	58.1	12.3	61.6	17.0	57.9	13.8	61.6	13.7	59.8	15.4	57.7
Dyna-Gro 9151	R	14.2	59.1	15.2	58.6	14.7	58.2	11.4	60.8	15.7	58.2	14.5	60.9	13.6	59.7	14.2	57.4
Dyna-Gro 9172	R	14.5	58.0	15.7	58.9	15.2	55.3	12.6	60.0	15.5	57.2	13.9	60.0	13.8	59.0	14.5	55.9
Dyna-Gro 9182	R	14.5	58.1	14.9	58.4	14.5	56.3	13.4	60.0	14.9	57.7	14.5	59.8	14.1	58.9	15.6	55.7
Dyna-Gro 9242W	W	14.7	57.9	14.6	58.3	15.2	55.8	13.1	59.8	16.4	57.5	13.7	59.2	13.9	58.3	15.5	56.6
Dyna-Gro WX19799W	W	14.4	56.3	14.4	58.3	14.7	53.3	13.1	57.0	16.6	55.5	13.6	58.0	13.7	57.8	14.8	54.3
Dyna-Gro WX20734	R	14.6	58.3	15.1	58.3	14.3	57.4	13.4	59.9	16.2	57.1	13.7	59.9	14.2	59.6	15.2	56.2
Dyna-Gro WX20738	R	14.5	58.5	14.5	59.1	14.5	57.3	12.6	60.0	16.7	57.2	13.4	60.8	14.2	59.3	15.3	56.2
Dyna-Gro WX21791W	W	14.6	58.2	14.8	58.4	14.9	54.9	13.0	60.1	16.5	57.0	14.2	60.3	14.1	59.7	14.8	57.1
Erisman	R	14.2	58.9	14.5	58.9	13.8	58.7	13.0	59.8	16.3	58.0	13.8	60.3	13.9	59.7	14.0	57.5
Harbor	R	14.4	57.9	14.5	58.5	14.9	56.7	12.8	59.4	16.1	57.6	13.5	59.1	13.6	58.3	15.1	56.0
Haubert	R	14.7	58.2	15.0	58.7	14.3	57.5	13.4	59.4	16.3	56.9	14.3	59.7	14.0	59.4	15.4	55.7
HS338R	R	14.5	58.7	14.7	58.7	14.4	58.3	12.7	59.2	16.9	57.3	14.2	60.0	13.9	60.5	14.8	57.1
ISF 1001	W	14.5	56.8	15.0	57.9	14.6	55.8	13.3	57.7	15.8	55.4	14.0	57.7	13.9	58.1	14.7	55.3
ISF 1115	W	14.2	57.3	14.4	58.0	14.3	57.1	11.7	57.5	16.6	56.0	14.2	58.6	13.9	58.9	14.0	55.3
ISF 790	R	14.6	58.0	16.8	58.2	14.1	56.8	12.2	58.8	15.0	56.3	14.1	60.6	14.0	59.7	15.8	55.4
Jupiter	W	14.4	57.8	15.3	57.9	14.8	55.9	13.2	60.1	15.1	57.3	13.7	59.0	13.7	57.9	14.9	56.4
KWS305	W	14.5	58.0	16.1	58.1	14.5	54.4	13.0	61.5	15.3	57.7	13.7	60.7	13.7	57.3	14.9	56.5
KWS308	W	15.0	57.7	17.6	58.2	14.9	56.2	12.6	58.2	17.0	57.6	14.7	59.1	13.7	58.4	14.3	56.1
KWS316	W	14.3	57.0	14.8	58.0	14.4	55.6	12.8	57.7	15.5	55.9	13.9	58.1	14.1	58.1	14.9	55.5
KWS317	W	14.3	57.5	14.3	58.3	14.4	56.8	13.4	58.3	15.7	56.3	14.0	58.7	14.0	59.0	14.4	55.3
KWS319	W	14.2	56.6	14.7	57.9	14.1	53.2	11.9	57.7	16.5	56.2	13.8	57.8	13.8	57.8	14.4	55.3
KWS327	W	14.2	56.4	16.2	58.5	14.4	53.2	12.1	56.5	14.9	56.7	13.6	58.1	13.9	57.4	14.3	54.5
KWS340	R	14.5	59.6	15.5	59.0	14.3	58.2	13.1	61.7	15.4	59.3	14.1	61.4	13.9	60.1	14.9	58.1
KWS356	R	14.1	58.5	14.6	58.6	13.7	56.9	13.4	59.1	14.8	58.0	13.4	59.4	13.9	59.8	14.7	57.3
KWS361	R	14.1	58.0	15.0	58.2	14.1	56.2	11.7	60.0	15.9	57.8	13.3	59.0	13.9	58.0	15.0	56.8

Table 2. Multi-Location Performance Summary for Test Weight and Percent Moisture.

		Ove	rall	Hilln	nan	Ingh	am	Hur	on	Sani	lac	Tusc	ola	Lenav	wee	Isabe	ella
Line	Color	% Moist	TW														
KWS375	R	14.3	58.9	15.8	58.8	14.4	57.2	12.1	59.5	15.1	59.2	13.8	60.7	13.9	60.7	14.9	56.7
LCS3334	R	14.6	58.8	15.1	58.5	14.7	58.4	11.9	59.3	17.2	57.5	14.3	60.2	13.9	59.9	14.6	57.8
LW2068	R	14.1	57.6	14.7	58.4	14.0	57.5	13.2	58.8	15.1	55.7	13.5	58.8	13.7	58.1	14.2	55.9
LW2148	R	14.5	57.9	14.8	58.4	14.9	55.6	12.4	59.9	16.8	56.5	13.4	59.1	13.9	58.7	14.9	57.0
LW2169	R	14.2	57.7	15.9	58.5	14.0	55.7	11.8	59.5	15.0	56.5	14.1	59.4	14.1	58.8	14.7	55.5
LW2958	R	14.3	58.4	14.4	58.5	13.9	56.4	12.8	60.6	16.5	57.2	13.5	60.1	14.1	59.5	15.1	56.9
MCIA Flipper	R	14.6	57.8	15.2	58.1	14.6	56.5	13.5	58.6	15.9	57.2	13.9	59.2	13.9	58.9	15.0	56.0
MCIA Jonah	R	14.8	57.7	16.5	57.9	14.5	56.8	13.1	58.6	16.5	57.4	13.9	59.2	14.1	58.7	14.9	55.6
MCIA MARLIN	R	14.7	58.3	18.0	57.6	13.3	56.4	12.9	59.8	15.9	56.9	13.8	61.2	14.4	59.8	14.8	56.2
MCIA Red Dragon	R	14.3	57.5	14.4	58.0	14.4	56.5	13.0	58.6	15.3	56.6	13.7	58.7	14.2	58.7	14.8	56.0
MCIA Whale	R	14.8	58.1	18.2	56.9	15.0	55.9	12.7	59.6	14.7	57.7	14.3	59.8	13.8	59.5	15.1	57.3
MCIA Wharf	R	13.8	56.9	14.2	58.3	13.7	54.5	12.8	59.2	14.4	56.7	13.4	57.7	14.0	56.7	14.1	55.6
MI14W0190	W	14.2	59.1	14.8	59.0	15.0	57.6	12.2	60.9	15.8	58.6	13.4	60.4	13.9	59.8	14.6	57.8
MI16R0720	R	14.3	56.5	14.7	58.0	13.8	55.4	13.5	57.3	15.5	55.3	13.8	58.0	13.9	57.3	14.7	54.0
MI16R0898	R	14.9	58.5	16.5	57.6	15.0	56.2	13.0	59.6	16.2	59.1	14.5	59.8	14.2	59.0	15.0	57.9
MI16R0906	R	14.2	57.7	15.4	57.9	14.6	55.5	12.6	58.6	14.9	57.2	13.8	59.3	13.6	58.6	14.3	56.4
MI16W0133	W	14.3	57.0	16.3	57.4	14.6	55.2	13.1	58.7	14.5	56.1	13.4	58.6	13.5	58.1	14.6	55.0
MI16W0528	W	14.3	57.0	14.4	58.3	14.5	54.4	13.4	58.7	15.3	56.4	14.5	58.0	14.1	58.1	13.9	55.3
MI17R0357	R	14.3	57.6	15.0	57.9	14.6	56.1	12.7	58.9	15.0	56.5	14.2	58.8	14.1	58.8	14.6	56.3
MI17W0224	W	14.9	57.4	17.4	57.7	14.3	55.5	13.5	58.7	15.3	57.6	14.4	58.6	14.1	58.0	15.5	56.2
MI17W0235	W	14.2	57.0	14.7	58.2	14.7	56.1	12.8	58.5	14.9	56.0	13.6	57.6	13.8	57.8	14.8	54.9
MI18R1605	R	14.1	58.2	14.4	58.5	14.0	57.4	12.6	59.6	15.8	56.6	13.8	60.2	13.5	59.4	14.0	56.3
MI18W0200	W	14.0	57.9	14.2	58.3	14.4	56.4	13.2	60.2	14.9	56.7	13.3	58.8	13.7	59.2	14.3	56.3
MI18W0286	W	14.3	56.5	14.5	58.0	15.3	53.2	12.1	59.2	15.8	54.2	13.6	58.6	13.7	58.2	14.9	54.2
Moonlight	W	14.3	56.8	14.5	58.0	14.3	54.6	12.9	58.5	15.3	56.0	14.2	57.5	13.8	57.3	14.6	56.0
Prestyn	R	14.1	57.4	14.4	58.3	14.5	56.9	12.7	58.1	15.2	56.8	13.6	58.8	14.0	58.4	14.4	54.8
REXP1215	R	14.2	57.2	14.1	58.2	13.7	56.0	12.9	59.6	16.3	55.9	13.3	58.7	14.1	57.1	14.8	55.2

Table 2. Multi-Location Performance Summary for Test Weight and Percent Moisture.

		Ove	rall	Hilln	nan	Ingh	am	Hur	on	San	ilac	Tusc	ola	Lenav	vee	Isabe	ella
Line	Color	% Moist	TW														
RS 902	R	14.1	58.4	14.9	58.4	13.8	58.3	11.9	58.9	15.4	57.6	14.0	59.9	14.0	59.8	14.8	56.0
RS 912	R	14.7	58.0	14.6	58.6	14.5	56.5	13.6	58.8	16.9	57.0	13.7	59.7	13.8	58.8	15.4	56.4
RS 961	R	14.4	57.6	15.0	58.7	14.6	56.9	13.2	59.3	16.4	57.2	13.5	59.6	14.3	58.9	14.1	52.1
RS 977	R	14.1	57.6	14.7	58.3	14.3	57.0	13.5	59.0	14.8	56.4	13.4	59.5	13.8	57.9	14.3	55.5
RWEXP1212	R	14.6	59.6	16.1	58.8	13.9	58.3	13.6	61.4	16.0	59.3	13.8	61.4	14.1	60.3	14.8	58.0
Sunburst	R	14.6	59.8	17.2	58.4	14.2	57.5	12.3	60.6	15.5	59.5	13.7	62.2	14.1	61.4	15.4	58.7
SY 100	R	14.2	56.3	16.2	57.8	13.7	55.2	12.3	57.2	15.3	55.8	13.4	57.5	14.1	57.4	14.4	53.6
SY 547	R	13.9	58.3	14.9	58.5	14.6	56.0	11.1	59.9	13.9	56.1	13.7	60.2	13.7	59.7	14.9	57.8
SY 576	R	14.0	57.8	15.0	58.7	14.8	54.5	11.5	59.2	14.8	57.4	13.4	60.4	13.8	58.5	14.8	55.7
SY Viper	R	14.4	59.0	16.0	58.1	13.4	58.3	11.8	58.4	16.4	58.8	13.8	61.1	14.0	60.5	15.5	57.9
Tyson	R	14.4	58.0	15.5	58.7	13.6	56.7	13.1	59.4	16.5	56.5	13.9	59.5	13.8	59.0	14.6	56.2
W 300	R	14.0	57.7	14.3	58.1	14.5	57.4	12.7	58.6	15.1	56.8	13.6	59.1	13.7	58.4	13.9	55.3
W 304	R	14.5	58.5	14.9	58.6	14.5	58.1	12.5	59.9	16.2	56.9	13.8	59.8	13.7	59.8	15.5	56.5
W 305	R	14.3	58.1	15.3	58.5	15.2	56.7	11.7	58.8	15.2	57.4	13.6	60.0	13.8	59.1	15.1	56.6
W 310	R	14.4	57.5	14.6	58.0	14.8	56.3	13.1	59.2	16.2	56.5	13.9	58.9	13.9	58.4	14.5	55.1
W 313	R	14.5	58.0	14.6	58.5	14.8	56.9	12.6	58.2	16.5	57.1	13.6	59.7	13.9	58.9	15.4	56.5
W 322	R	14.1	58.3	14.9	58.7	13.5	57.1	13.0	60.2	14.8	57.1	13.3	60.2	13.8	59.6	14.8	55.7
W 324	R	14.4	57.9	15.6	58.4	14.3	57.2	12.6	59.3	15.3	56.5	13.9	59.6	14.0	58.6	14.7	56.1
Whitetail	W	14.0	57.3	14.4	58.2	14.7	55.7	12.9	59.3	14.3	55.7	13.5	58.1	14.1	58.3	14.4	55.7
	Mean	14.3	57.9	15.1	58.3	14.4	56.3	12.8	59.3	15.6	57.0	13.8	59.4	13.9	58.8	14.8	56.1
	CV	1.7	1.4	6.4	1.0	3.3	2.3	4.3	1.8	4.6	1.7	2.4	1.9	1.4	1.6	2.8	1.9
	LSD	0.3	0.4	0.6	0.5	1.3	8.0	1.7	8.0	1.7	1.1	0.6	1.3	0.4	0.4	0.8	1.5

Table 3. Fusarium Head Blight Resistance, lodging, cephalosporium stripe, pre-harvest sprouting, plant height and flowering data.

Table 3. Fusarium Head	Jugue neon	l loug		ium Head		cot oproutii	ig, plant nei	gine uniu novi	reinig autur	Preharvest	Plant	Flowering	Pysiological	Grain Fill
		Severity	Incidence	Index	DON ppm	FHB	Lodging	Cephalospo	orium Stripe	Sprouting	Height	Date	Maturity	Period
Line	Color	2021	2021	2021	2020	Rating*	(0-9)**	Severity	Incidence	(0-9)**	(inches)	Days past Jan. 1	Days past Jan. 1	# of days
13VTK59-55	R	49.7	94.8	47.1			1	2.1	20.2	2.6	28.5	146	171	25
AC Mountain	W	47.6	94.6	45.0	43.9		3.5	2.4	17.5	6.7	37.4	147	175	28
AgriMAXX 498	R	42.1	94.6	39.8	25.9		2.5	2.1	30.8	2.9	34.6	147	178	31
AgriMAXX 502	R	42.4	94.8	40.2			2	2.3	46.5	2.2	31.2	146	175	29
AgriMAXX 503	R	40.6	94.6	38.5	11.9		5	2.8	49.7	1.4	33.7	147	175	28
AgriMAXX 505	R	46.7	94.7	44.2	13		1.5	2.1	61.8	1.3	30.3	148	175	27
AgriMAXX 513	R	39.1	93.5	36.6			5	2.3	53.9	0.8	32.7	147	175	28
AgriMAXX 516	R	45.1	94.5	42.6			0	1.6	78.4	0.0	31.8	147	177	30
AgriMAXX EXP 2002	R	37.6	94.0	35.3			1	2.1	64.1	1.3	30.3	146	176	30
AgriMAXX EXP 2055W	W	45.1	94.6	42.6	38.5		4.5	2.0	64.2	6.0	32.6	146	176	30
AgriMAXX Exp. 2050W	W	43.6	94.6	41.3	22.8		2.5	1.9	25.9	6.5	30.1	147	177	30
Ambassador	W	73.5	94.7	69.6	51.9		3	2.1	49.6	5.4	35.7	147	175	28
DF 105 R	R	49.6	94.6	46.9	25.4		3.5	2.3	17.2	4.2	30.6	146	175	29
DF 112 R	R	54.1	94.7	51.3	18.8		1.5	1.8	49.7	2.0	30.9	146	175	29
DF 119 R	R	49.6	94.8	47.0	18.7		2.5	2.7	46.5	2.1	31.4	146	169	23
DF 121 R	R	54.1	94.8	51.3			1	1.8	75.9	0.0	31.9	147	176	29
DF 131 R	R	43.6	94.6	41.2			2	2.2	43.4	3.2	30.9	146	177	31
DF 141 R	R	55.7	94.7	52.8	16.7		0	1.9	82.3	3.7	29.3	146	175	29
DF 218 W	W	56.0	94.6	53.0	55.4		0.5	2.7	14.1	5.6	31.7	148	176	28
DF 261 W	W	50.3	94.5	47.5			4	2.0	43.5	3.4	30.8	146	171	25
DF 271 W	W	45.2	94.8	42.9			1	2.4	27.7	2.8	30.4	147	173	26
DF EX 2101 R	R	61.2	94.6	57.9			1	1.9	56.2	0.0	31.5	147	174	27
DF EX 2102 R	R	33.2	93.0	30.9			1.5	2.0	51.6	0.9	31.5	146	174	28
DF EX 2103 R	R	49.7	94.8	47.2			5.5	2.2	26.2	0.8	30.8	146	173	27
DF EX 2104 R	R	41.5	94.6	39.2			5	2.4	91.8	4.8	32.6	148	173	25
Dyna-Gro 9002	R	49.4	94.6	46.7	16.6		4	1.8	58.4	2.8	32.1	146	174	28
Dyna-Gro 9070	R	36.2	94.2	34.1	30.2		2	1.8	62.4	3.7	31.7	146	175	29
Dyna-Gro 9082W	W	34.6	94.5	32.7	26.2		0	1.9	70.6	6.2	30.7	148	177	29
Dyna-Gro 9120	R	31.6	92.6	29.3			0	1.8	37.3	1.5	30.2	146	177	31
Dyna-Gro 9151	R	42.2	94.6	39.9	26.8		2	2.2	37.2	2.2	31.1	148	175	27

Table 3. Fusarium Head Blight Resistance, lodging, cephalosporium stripe, pre-harvest sprouting, plant height and flowering data.

Table 3. Fusarium Head	Diigitt Kesis	l loug		ium Head		est sproutii	ig, plant nei	giit and now	vering data.	Preharvest	Plant	Flowering	Pysiological	Grain Fill
		Severity	Incidence	Index	DON ppm	FHB	Lodging	Cephalospo	orium Stripe		Height	Date	Maturity	Period
Line	Color	2021	2021	2021	2020	Rating*	(0-9)**	Severity	Incidence	(0-9)**	(inches)	Days past Jan. 1	Days past Jan. 1	# of days
Dyna-Gro 9172	R	45.5	94.8	43.2	11.8		1	2.5	67.9	1.3	31.4	147	177	30
Dyna-Gro 9182	R	43.6	94.7	41.3	19.3		2	2.3	52.7	1.1	33	147	175	28
Dyna-Gro 9242W	W	33.1	94.6	31.3	47.8		0	2.3	58.2	4.2	32.9	147	176	29
Dyna-Gro WX19799W	W	51.5	94.6	48.8			2.5	1.9	60.3	6.3	30.3	147	176	29
Dyna-Gro WX20734	R	51.9	94.6	49.1			2	1.8	70.9	2.0	30.3	146	174	28
Dyna-Gro WX20738	R	42.1	94.7	39.9			3	2.5	61.6	1.7	32	146	175	29
Dyna-Gro WX21791W	W	39.1	94.2	36.8			0	1.9	65.4	6.8	30	147	176	29
Erisman	R	37.4	94.5	35.3	8.5		8	2.7	31.7	2.6	33.3	147	169	22
Harbor	R	45.2	94.7	42.8	30.8		7.5	2.3	39.7	1.6	32.7	146	172	26
Haubert	R	40.7	94.7	38.5			2	2.8	37.7	0.2	32.4	146	175	29
HS338R	R	37.7	94.8	35.7	10.6		8	2.0	44.0	0.8	32.3	146	173	27
ISF 1001	W	46.6	94.7	44.2			5	2.2	71.8	5.8	31.5	146	174	28
ISF 1115	W	37.6	94.5	35.5			2.5	2.9	35.9	4.3	28.9	146	173	27
ISF 790	R	46.7	94.7	44.2			7	1.6	44.3	3.8	31.5	146	174	28
Jupiter	W	66.0	94.5	62.4	35.8		2	2.4	37.1	4.1	30.8	147	177	30
KWS305	W	43.6	94.6	41.3			8	2.6	34.8	3.3	32.4	147	176	29
KWS308	W	42.1	94.5	39.8			8.5	2.6	52.0	2.2	31.7	147	175	28
KWS316	W	53.5	94.8	50.8	18		3	2.5	47.7	6.4	30.7	146	174	28
KWS317	W	43.7	94.5	41.3	24.4		3	2.1	42.2	4.7	30.8	146	174	28
KWS319	W	46.7	94.8	44.3			4	1.8	62.1	4.6	31	147	172	25
KWS327	W	61.6	94.9	58.5			4.5	2.3	45.9	7.0	29.4	147	173	26
KWS340	R	46.6	93.6	43.7			3	2.1	42.2	0.2	31.9	146	175	29
KWS356	R	57.2	94.7	54.2			0	2.4	71.0	6.5	30.7	148	177	29
KWS361	R	39.1	94.7	37.1			4.5	2.2	68.0	0.8	35	147	173	26
KWS375	R	42.2	94.8	40.0			2.5	2.1	24.2	1.4	29.3	147	176	29
LCS3334	R	31.7	93.8	29.7	18.1		6	2.2	14.5	0.2	33.6	146	175	29
LW2068	R	33.3	94.8	31.5	19.4		3	2.8	13.6	1.7	31.2	146	175	29
LW2148	R	34.6	94.5	32.7			6.5	3.0	70.4	0.0	34.3	147	174	27
LW2169	R	41.1	94.7	38.9			0	2.6	43.6	1.4	30	147	173	26
LW2958	R	34.6	94.5	32.7			2	2.2	50.8	1.4	32.4	146	171	25

2021 Michigan State University Wheat Performance Trials Table 3. Fusarium Head Blight Resistance, lodging, cephalosporium stripe, pre-harvest sprouting, plant height and flowering data.

Table 3. Fusarium Head	Blight Kesis	tance, loug		porium str ium Head		est sproutin	ig, plant nei	gnt and flow	ering data.	Preharvest	Plant	Flowering	Pysiological	Grain Fill
		Severity	Incidence	Index	DON ppm	FHB	Lodging	Cephalospo	rium Stripe	Sprouting	Height	Date	Maturity	Period
Line	Color	2021	2021	2021	2020	Rating*	(0-9)**	Severity	Incidence	(0-9)**	(inches)	Days past Jan. 1	Days past Jan. 1	# of days
MCIA Flipper	R	52.3	94.8	49.6	22.1		0.5	2.0	36.3	1.3	28.9	146	173	27
MCIA Jonah	R	51.1	94.6	48.3	24.3		2.5	2.3	49.2	1.4	32.2	146	176	30
MCIA MARLIN	R	61.4	94.6	58.1	25.8		7	1.7	42.7	2.4	28	147	174	27
MCIA Red Dragon	R	49.7	92.7	46.0	19.8		3	2.3	52.4	1.5	36.9	146	172	26
MCIA Whale	R	47.7	94.5	45.1	23.8		0.5	2.9	11.0	3.8	34.6	148	176	28
MCIA Wharf	R	48.2	94.3	45.5	9.5		2	2.7	27.1	4.0	27.4	146	175	29
MI14W0190	W	19.8	94.6	18.7	23.9		0.5	2.6	33.0	4.6	32.3	146	172	26
MI16R0720	R	60.1	94.7	56.9	25		6	2.0	52.6	2.5	29.5	147	171	24
MI16R0898	R	26.8	94.5	25.3	17.5		1	2.4	28.9	0.0	36.3	148	178	30
MI16R0906	R	58.5	94.6	55.4	23.5		1	1.9	37.1	3.1	31.5	146	177	31
MI16W0133	W	58.6	94.7	55.5	56.6		2	2.0	38.9	6.3	31.1	146	174	28
MI16W0528	W	63.0	94.8	59.7	35.4		4.5	2.2	27.2	4.3	32.3	147	176	29
MI17R0357	R	55.6	94.7	52.7	27.3		1	1.9	64.7	1.5	27.4	146	174	28
MI17W0224	W	60.0	94.5	56.7	54		3	2.4	11.9	5.5	29.9	146	174	28
MI17W0235	W	64.6	94.7	61.2	29.8		5	2.2	43.9	4.6	34	147	176	29
MI18R1605	R	52.0	94.3	49.0			0.5	1.5	38.1	2.0	30.9	146	169	23
MI18W0200	W	73.6	95.0	69.9			6	2.4	43.7	3.7	29.4	147	171	24
MI18W0286	W	65.0	94.6	61.5			0	1.8	61.4	5.6	30.9	147	178	31
Moonlight	W	66.1	94.6	62.5	34.1		7	2.4	37.7	2.3	32.2	146	173	27
Prestyn	R	33.1	94.7	31.3			7.5	2.4	55.1	0.8	32.2	146	173	27
REXP1215	R	40.7	94.7	38.5			3	2.2	49.4	2.1	29	146	172	26
RS 902	R	39.1	94.6	37.0	15.1		0.5	2.1	24.0	0.2	31.2	146	175	29
RS 912	R	37.2	94.6	35.1	9.8		4	2.4	58.7	0.8	33.8	148	172	24
RS 961	R	45.5	94.8	43.1	22.8		1.5	2.4	54.4	0.6	30	147	176	29
RS 977	R	39.3	94.8	37.2	13.5		3.5	1.8	51.2	1.5	30.9	147	176	29
RWEXP1212	R	41.4	94.8	39.2			4	1.8	78.0	0.8	30.2	146	173	27
Sunburst	R	45.8	94.7	43.4	20.4		1	2.5	40.0	0.0	28.5	148	179	31
SY 100	R	58.6	94.7	55.5	27.7		4	2.3	37.7	2.7	30.7	146	177	31
SY 547	R	37.7	94.2	35.5	33.8		2	1.1	70.4	1.3	33	146	173	27
SY 576	R	43.6	94.6	41.2	15.4		3	2.4	65.3	3.7	35	149	177	28

Table 3. Fusarium Head Blight Resistance, lodging, cephalosporium stripe, pre-harvest sprouting, plant height and flowering data.

			Fusar	ium Head	Blight					Preharvest	Plant	Flowering	Pysiological	Grain Fill
		Severity	Incidence	Index	DON ppm	FHB	Lodging	Cephalospo	orium Stripe	Sprouting	Height	Date	Maturity	Period
Line	Color	2021	2021	2021	2020	Rating*	(0-9)**	Severity	Incidence	(0-9)**	(inches)	Days past Jan. 1	Days past Jan. 1	# of days
SY Viper	R	42.2	93.6	39.5	22.1		5	2.4	50.2	1.6	35.1	146	172	26
Tyson	R	41.6	94.7	39.4			0.5	2.0	33.3	0.8	29.6	146	173	27
W 300	R	36.1	94.5	34.2			1	2.5	30.3	0.8	31.8	146	173	27
W 304	R	37.7	94.6	35.6	17.3		1	3.2	0.0	0.3	32.8	146	176	30
W 305	R	40.7	94.8	38.6	11.3		2.5	2.8	19.2	1.2	31.3	147	177	30
W 310	R	40.7	94.8	38.6	19.2		2	2.4	39.2	2.4	30.6	147	176	29
W 313	R	34.6	94.6	32.7	12.1		5	2.1	90.3	0.8	32.4	146	173	27
W 322	R	42.1	94.8	39.9			0	1.9	49.7	0.6	32.8	147	175	28
W 324	R	43.2	94.7	40.9			2	1.9	51.7	1.3	30	147	173	26
Whitetail	W	49.7	94.4	47.0			5	1.5	61.5	6.8	31.8	146	173	27
	Mean	46.1	94.5				2.9	2.2	46.6	2.7	31.6	146.6	174.5	27.9
	CV	21.5	0.4				76.8	16.3	39.4	75.5	6.0	0.5	1.2	
	LSD	3.2	0.7				3.8	0.7	34.5	3.2	2.3	1.0	3.0	

Table 4. Conventional (00, 13		Fuscola	.,		Tuscola		Tusco	ola		Isabella			Isabella		Isab	ella
		High N	/lanageme	nt	Conventio	nal Manag	gement	нм - с	onv.	High	Manager	nent	Conver	ntional Ma	nagement	нм - с	Conv.
Line	Color	Bu/A	% Moist	TW	Bu/A	% Moist	TW	Difference	Rank	Bu/A	% Moist	TW	Bu/A	% Moist	TW	Difference	Rank
13VTK59-55	R	91.4	13.7	61.3	83.8	14.6	60.4	7.6	42	103.6	14.9	57.3	88.2	15.4	56.1	15.4	3
AC Mountain	W	86.8	13.3	57.8	89.5	14.1	57.6	-2.8	96	96.5	14.8	55.0	91.5	15.0	54.2	5.0	81
AgriMAXX 498	R	96.2	14.4	59.0	93.6	14.5	58.9	2.6	74	105.3	14.9	55.4	97.4	16.0	53.9	7.9	54
AgriMAXX 502	R	101.4	13.9	59.6	88.6	14.1	59.0	12.8	19	103.4	15.0	55.4	99.7	15.1	54.4	3.7	92
AgriMAXX 503	R	105.7	14.2	60.2	93.6	14.3	58.9	12.1	21	107.6	14.6	56.4	99.6	15.3	55.2	8.0	51
AgriMAXX 505	R	101.9	13.6	61.4	100.4	13.7	61.2	1.5	84	104.2	14.6	57.1	94.0	14.6	55.4	10.2	30
AgriMAXX 513	R	97.3	13.7	60.7	95.0	14.1	59.5	2.3	76	104.6	14.5	56.7	95.8	15.1	55.3	8.8	39
AgriMAXX 516	R	100.0	14.1	59.3	89.0	14.1	58.4	11.0	24	107.8	14.3	55.8	99.8	15.0	54.8	8.0	50
AgriMAXX EXP 2002	R	95.1	14.0	60.8	83.2	14.1	60.6	12.0	22	103.6	15.1	57.5	93.4	15.4	55.9	10.3	27
AgriMAXX EXP 2055W	W	95.3	14.0	58.0	93.3	14.2	57.6	2.1	80	97.1	14.9	55.9	91.5	15.2	54.5	5.6	76
AgriMAXX Exp. 2050W	W	114.7	13.6	58.3	95.1	14.2	58.2	19.6	2	109.2	14.3	55.5	100.6	14.5	54.7	8.7	41
Ambassador	W	87.1	13.5	57.0	83.6	13.5	56.4	3.4	68	100.5	14.3	54.9	94.9	14.6	54.2	5.7	75
DF 105 R	R	97.5	13.7	58.8	93.6	14.1	58.1	3.9	64	101.7	14.7	55.6	93.1	15.0	54.1	8.6	42
DF 112 R	R	97.0	14.0	58.7	94.8	14.2	58.5	2.2	77	105.5	14.3	55.2	94.2	14.8	53.8	11.3	17
DF 119 R	R	97.9	13.9	59.6	94.4	14.0	59.0	3.5	65	104.0	15.1	56.7	96.9	15.3	55.7	7.1	59
DF 121 R	R	107.3	13.0	56.3	100.0	14.0	59.0	7.3	44	110.4	15.5	56.7	106.0	15.1	55.1	4.4	86
DF 131 R	R	101.2	13.7	59.7	96.6	14.4	58.6	4.6	61	106.2	14.6	56.8	97.8	14.6	54.7	8.5	44
DF 141 R	R	90.2	13.2	60.0	82.7	14.0	58.9	7.5	43	102.4	14.5	55.8	95.4	15.2	54.5	7.0	60
DF 218 W	W	95.7	14.0	60.0	88.5	14.2	58.2	7.2	45	100.9	15.2	55.7	98.5	15.5	56.0	2.4	96
DF 261 W	W	97.5	13.9	58.2	92.8	14.0	58.0	4.7	59	98.7	15.1	55.5	90.1	15.4	55.0	8.5	43
DF 271 W	W	111.4	13.4	59.4	95.3	14.3	58.0	16.1	6	108.1	14.3	55.0	101.2	14.8	54.0	7.0	61
DF EX 2101 R	R	100.4	13.5	60.7	92.2	14.2	59.8	8.3	38	105.3	14.8	55.7	98.9	14.6	54.5	6.4	67
DF EX 2102 R	R	97.3	13.7	60.1	83.1	14.2	59.5	14.2	13	105.3	15.2	57.2	92.7	15.5	56.1	12.6	12
DF EX 2103 R	R	97.5	14.0	57.1	92.1	14.0	56.8	5.3	57	110.3	14.3	53.9	99.2	14.6	52.8	11.1	21
DF EX 2104 R	R	99.0	13.8	61.0	93.9	14.3	60.5	5.1	58	103.6	14.6	58.3	96.7	14.7	57.3	7.0	62
Dyna-Gro 9002	R	96.5	14.2	59.1	102.3	14.7	58.6	-5.8	98	105.8	14.5	55.6	98.9	15.6	54.7	6.9	63
Dyna-Gro 9070	R	106.9	13.8	60.1	87.0	14.0	58.3	19.8	1	107.6	15.1	55.6	96.7	15.9	53.9	10.9	25
Dyna-Gro 9082W	W	89.9	13.6	58.7	90.1	13.7	58.3	-0.2	91	106.1	14.7	56.6	94.8	14.9	54.6	11.2	18
Dyna-Gro 9120	R	91.5	13.8	61.6	90.0	14.0	61.3	1.5	83	106.3	15.4	57.7	99.9	15.4	56.5	6.3	68
Dyna-Gro 9151	R	100.6	14.5	60.9	86.9	13.9	60.7	13.7	14	105.4	14.2	57.4	95.4	15.2	55.8	10.0	32
Dyna-Gro 9172	R	102.4	13.9	60.0	91.4	14.0	59.0	11.0	23	107.2	14.5	55.9	99.6	15.4	54.6	7.6	57

Table 4. Conventional (Tuscola	,		Гuscolа		Tusco	la		Isabella		Isabella			Isabe	lla
		High N	Nanageme	nt	Conventio	nal Mana	gement	HM - Co	onv.	High Management		Conventional Management			HM - C	onv.	
Line	Color	Bu/A	% Moist	TW	Bu/A	% Moist	TW	Difference	Rank	Bu/A	% Moist	TW	Bu/A	% Moist	TW	Difference	Rank
Dyna-Gro 9182	R	94.5	14.5	59.8	88.5	14.1	58.5	5.9	54	104.5	15.6	55.7	100.4	15.5	55.4	4.1	91
Dyna-Gro 9242W	W	90.2	13.7	59.2	97.9	14.3	58.4	-7.7	100	103.2	15.5	56.6	94.5	15.8	55.8	8.7	40
Dyna-Gro WX19799W	W	90.3	13.6	58.0	88.9	13.8	57.7	1.4	85	100.9	14.8	54.3	94.7	15.0	52.9	6.3	70
Dyna-Gro WX20734	R	100.7	13.7	59.9	92.7	14.5	58.7	8.0	41	105.4	15.2	56.2	101.0	16.0	54.9	4.4	85
Dyna-Gro WX20738	R	92.8	13.4	60.8	90.2	13.5	60.1	2.6	73	103.6	15.3	56.2	95.6	15.5	55.0	7.9	53
Dyna-Gro WX21791W	W	94.7	14.2	60.3	86.7	14.1	59.2	8.1	39	104.1	14.8	57.1	94.4	15.3	56.2	9.6	36
Erisman	R	88.0	13.8	60.3	77.2	14.0	59.1	10.7	26	91.0	14.0	57.5	87.7	15.3	56.2	3.3	94
Harbor	R	87.5	13.5	59.1	84.7	13.9	58.5	2.8	71	91.9	15.1	56.0	91.0	16.0	55.2	0.9	97
Haubert	R	98.6	14.3	59.7	89.0	14.1	59.4	9.6	32	104.4	15.4	55.7	99.4	15.5	55.0	5.0	79
HS338R	R	97.2	14.2	60.0	94.5	14.3	59.5	2.7	72	104.7	14.8	57.1	100.5	15.7	55.9	4.2	90
ISF 1001	W	93.4	14.0	57.7	91.4	14.1	57.6	2.0	81	100.2	14.7	55.3	87.1	15.2	54.7	13.1	10
ISF 1115	W	102.0	14.2	58.6	101.0	14.4	58.2	1.0	88	107.5	14.0	55.3	93.9	14.7	54.0	13.7	8
ISF 790	R	104.6	14.1	60.6	90.1	14.1	59.9	14.5	12	109.7	15.8	55.4	98.6	15.6	55.3	11.1	22
Jupiter	W	100.5	13.7	59.0	93.8	14.0	58.2	6.7	50	106.7	14.9	56.4	98.4	15.3	55.5	8.2	47
KWS305	W	91.6	13.7	60.7	83.6	13.8	59.3	8.0	40	96.8	14.9	56.5	88.9	15.2	55.5	7.8	55
KWS308	W	101.4	14.7	59.1	93.1	14.6	59.6	8.3	37	97.5	14.3	56.1	84.2	15.0	55.0	13.4	9
KWS316	W	92.3	13.9	58.1	85.4	14.5	57.4	6.9	49	95.8	14.9	55.5	87.6	15.5	54.3	8.2	46
KWS317	W	99.0	14.0	58.7	106.0	14.3	58.1	-6.9	99	104.4	14.4	55.3	99.9	14.8	54.2	4.5	83
KWS319	W	96.7	13.8	57.8	90.9	13.8	57.4	5.8	55	101.3	14.4	55.3	97.0	14.7	54.8	4.3	88
KWS327	W	97.7	13.6	58.1	93.1	13.9	57.2	4.6	60	92.3	14.3	54.5	86.6	14.6	53.6	5.7	74
KWS340	R	93.2	14.1	61.4	94.7	14.5	60.7	-1.6	95	106.1	14.9	58.1	98.1	15.8	56.5	8.1	49
KWS356	R	92.6	13.4	59.4	86.1	14.0	58.8	6.5	52	100.8	14.7	57.3	96.4	15.1	54.8	4.4	84
KWS361	R	107.7	13.3	59.0	92.6	14.2	58.7	15.1	7	105.3	15.0	56.8	100.0	15.6	55.5	5.3	77
KWS375	R	87.3	13.8	60.7	84.1	13.9	60.6	3.2	70	104.3	14.9	56.7	92.1	15.3	55.1	12.2	13
LCS3334	R	96.2	14.3	60.2	94.3	14.9	59.5	1.9	82	104.0	14.6	57.8	97.3	15.5	56.6	6.7	64
LW2068	R	102.7	13.5	58.8	93.3	14.0	58.3	9.4	33	107.7	14.2	55.9	101.9	14.5	54.5	5.8	72
LW2148	R	99.9	13.4	59.1	92.7	14.2	59.3	7.2	46	103.0	14.9	57.0	96.4	15.4	55.0	6.5	66
LW2169	R	98.6	14.1	59.4	99.8	14.3	58.8	-1.1	94	108.6	14.7	55.5	100.3	15.0	54.5	8.3	45
LW2958	R	95.1	13.5	60.1	90.8	13.8	59.0	4.3	63	103.0	15.1	56.9	93.5	15.2	55.9	9.4	37
MCIA Flipper	R	99.6	13.9	59.2	99.5	14.0	59.4	0.1	90	108.6	15.0	56.0	102.9	15.4	55.0	5.7	73
MCIA Jonah	R	91.7	13.9	59.2	89.4	14.4	58.8	2.4	75	103.4	14.9	55.6	99.1	15.5	54.6	4.4	87

Table 4. Conventional (Conv.) vs High Management (HM) Yield Results. Tuscola Tuscola Tuscola Isabella								lach - U		Isabella						
			uscola			Fuscola		HM - Conv.		11:04	Isabella		Isabella Conventional Management			HM - Conv.	
Line	Color	Bu/A	lanageme % Moist	TW	Conventio Bu/A	nai iviana _§ % Moist	gement TW	Difference	Rank	Bu/A	Managen % Moist	TW	Bu/A	% Moist	TW	Difference	Rank
MCIA MARLIN	R	96.3	13.8	61.2	97.2	14.3	59.8	-1.0	93	108.2	14.8	56.2	105.4	16.0	55.2	2.8	95
MCIA Red Dragon	R	101.1	13.7	58.7	87.5	14.4	57.7	13.5	16	107.7	14.8	56.0	93.6	15.1	55.2	14.1	7
MCIA Whale	R	86.7	14.3	59.8	87.2	14.2	59.9	-0.6	92	104.4	15.1	57.3	86.2	15.9	55.3	18.2	1
MCIA Wharf	R	105.3	13.4	57.7	96.6	13.7	57.2	8.7	36	108.8	14.1	55.6	93.7	14.5	54.2	15.0	4
MI14W0190	W	90.3	13.4	60.4	80.1	14.4	59.1	10.2	29	100.5	14.6	57.8	85.7	15.8	55.9	14.8	6
MI16R0720	R	103.4	13.8	58.0	86.4	14.3	56.9	17.0	5	101.0	14.7	54.0	101.0	15.1	53.0	0.1	98
MI16R0898	R	93.0	14.5	59.8	89.5	14.4	59.4	3.5	65	103.8	15.0	57.9	92.7	15.5	56.5	11.1	23
MI16R0906	R	103.6	13.8	59.3	89.9	14.0	57.7	13.7	14	104.8	14.3	56.4	94.7	14.7	55.2	10.1	31
MI16W0133	W	92.5	13.4	58.6	89.0	14.2	57.5	3.5	67	103.3	14.6	55.0	95.7	15.1	54.0	7.6	57
MI16W0528	W	97.7	14.5	58.0	97.2	14.2	57.6	0.5	89	100.1	13.9	55.3	90.4	14.6	54.0	9.7	35
MI17R0357	R	103.1	14.2	58.8	88.2	13.9	58.4	15.0	8	106.7	14.6	56.3	97.7	15.3	54.6	9.0	38
MI17W0224	W	102.1	14.4	58.6	91.4	14.5	58.2	10.7	27	98.2	15.5	56.2	93.9	15.5	55.4	4.3	88
MI17W0235	W	100.9	13.6	57.6	86.1	14.3	57.2	14.8	10	106.6	14.8	54.9	96.3	14.8	54.3	10.3	26
MI18R1605	R	99.7	13.8	60.2	88.9	12.9	56.8	10.8	25	104.0	14.0	56.3	92.3	15.1	54.7	11.7	14
MI18W0200	W	95.0	13.3	58.8	77.8	13.8	56.7	17.2	4	101.9	14.3	56.3	87.0	14.2	55.3	14.9	5
MI18W0286	W	98.3	13.6	58.6	92.8	14.2	56.9	5.5	56	100.4	14.9	54.2	87.8	15.3	52.7	12.6	11
Moonlight	W	97.0	14.2	57.5	84.7	14.5	56.7	12.3	20	98.8	14.6	56.0	90.7	15.2	54.7	8.1	48
Prestyn	R	90.7	13.6	58.8	89.6	13.7	58.3	1.1	86	100.9	14.4	54.8	89.6	14.9	54.2	11.4	16
REXP1215	R	108.0	13.3	58.7	93.4	14.4	57.7	14.6	11	103.9	14.8	55.2	105.7	15.1	53.9	-1.8	100
RS 902	R	96.4	14.0	59.9	87.3	14.0	59.5	9.1	34	107.4	14.8	56.0	96.1	15.6	55.0	11.2	18
RS 912	R	104.1	13.7	59.7	84.5	14.2	59.0	19.6	3	103.6	15.4	56.4	95.6	15.6	54.7	7.9	52
RS 961	R	85.0	13.5	59.6	82.9	14.2	59.1	2.1	79	109.5	14.1	52.1	93.3	14.9	55.3	16.2	2
RS 977	R	94.6	13.4	59.5	97.5	14.3	58.3	-2.9	97	103.8	14.3	55.5	97.1	15.4	54.0	6.7	65
RWEXP1212	R	100.2	13.8	61.4	93.2	13.9	60.9	6.9	48	107.8	14.8	58.0	104.4	15.9	56.6	3.4	93
Sunburst	R	96.3	13.7	62.2	89.2	14.0	61.4	7.1	47	103.5	15.4	58.7	97.2	15.8	58.2	6.3	69
SY 100	R	106.0	13.4	57.5	99.3	14.3	56.6	6.7	51	105.0	14.4	53.6	100.0	15.5	52.0	5.0	80
SY 547	R	101.6	13.7	60.2	97.2	14.2	59.3	4.4	62	108.4	14.9	57.8	98.2	15.5	56.0	10.2	28
SY 576	R	91.0	13.4	60.4	87.7	13.7	59.8	3.3	69	100.3	14.8	55.7	92.6	15.0	54.9	7.7	56
SY Viper	R	90.5	13.8	61.1	88.3	14.5	60.1	2.2	77	106.4	15.5	57.9	100.6	16.2	56.9	5.8	71
Tyson	R	100.0	13.9	59.5	93.8	14.1	58.6	6.2	53	108.6	14.6	56.2	98.3	15.1	54.2	10.2	29

		Tuscola			Tuscola			Tuscola		Isabella			Isabella			Isabella	
		High N	/lanageme	nt	Conventional Management			HM - Conv.		High Management			Conventional Management			HM - Conv.	
Line	Color	Bu/A	% Moist	TW	Bu/A	% Moist	TW	Difference	Rank	Bu/A	% Moist	TW	Bu/A	% Moist	TW	Difference	Rank
W 300	R	94.3	13.6	59.1	84.7	14.2	58.1	9.6	31	92.6	13.9	55.3	93.6	15.3	53.9	-1.1	99
W 304	R	93.4	13.8	59.8	92.4	14.4	59.2	1.0	87	106.3	15.5	56.5	96.3	15.6	54.8	10.0	33
W 305	R	101.8	13.6	60.0	86.8	14.4	58.9	14.9	9	105.1	15.1	56.6	93.7	15.4	54.8	11.5	15
W 310	R	103.8	13.9	58.9	94.0	14.1	58.3	9.9	30	102.3	14.5	55.1	97.2	14.8	54.8	5.1	78
W 313	R	101.6	13.6	59.7	88.1	14.2	58.9	13.4	17	100.8	15.4	56.5	96.1	15.4	54.7	4.7	82
W 322	R	102.8	13.3	60.2	92.3	13.8	59.7	10.6	28	107.9	14.8	55.7	98.1	15.0	54.5	9.9	34
W 324	R	102.7	13.9	59.6	89.6	14.3	58.5	13.2	18	112.4	14.7	56.1	101.4	15.1	54.7	11.1	24
Whitetail	W	96.9	13.5	58.1	88.0	14.2	57.0	8.9	35	104.0	14.4	55.7	92.9	14.8	54.7	11.2	20
	Mean	97.6	13.8	59.4	90.8	14.1	58.7	6.7		103.8	14.8	56.1	95.7	15.2	54.9	8.1	
	cv	5.9	2.4	1.9	5.8	2.0	1.9			4.0	2.8	1.9	4.8	2.7	1.8		
	LSD	9.3	0.6	1.3	7.2	0.9	1.2			2.3	1.5	0.8	3.8	0.8	0.9		

Table 5. Milling and baking qualities.

Table 5. Milling and bal	I	NIR Kernel		Adjusted Flour		Flour		Sodium	Cookie
		Protein	SKCS Kernel	Yield	Softness	Protein	Lactic Acid	Carbonate	Diameter
Line	Color	(at 12%)	Hardness	(%)	Equivalent (%)	(at 14%)	SRC (%)	SRC (%)	(cm)
13VTK59-55	R								
AC Mountain	W	9.9	26.3	69.0	55.1	7.9	83.5	66.3	19.3
AgriMAXX 498	R	9.6	26.5	70.2	61.2	7.7	104.1	66.7	18.9
AgriMAXX 502	R								
AgriMAXX 503	R	10.2	32.7	66.9	53.9	8.4	111.0	69.0	18.4
AgriMAXX 505	R	10.1	26.6	64.3	58.1	8.3	137.5	77.7	18.5
AgriMAXX 513	R								
AgriMAXX 516	R								
AgriMAXX EXP 2002	R								
AgriMAXX EXP 2055W	W	9.9	29.2	68.2	52.2	8.3	110.3	69.6	18.9
AgriMAXX Exp. 2050W	W	9.4	19.6	69.5	59.4	7.4	102.7	68.2	19.1
Ambassador	W	10.2	15.9	70.0	56.1	7.9	96.8	65.5	19.1
DF 105 R	R	9.4	38.6	68.9	54.9	7.8	93.0	67.7	19.4
DF 112 R	R	9.1	21.6	70.3	57.4	7.3	113.3	71.0	19.3
DF 119 R	R	10.1	20.7	68.6	58.5	7.9	108.6	67.7	19.8
DF 121 R	R								
DF 131 R	R								
DF 141 R	R	9.3	26.2	67.4	58.2	7.3	86.7	70.3	19.4
DF 218 W	w	10.0	29.4	67.8	51.6	8.3	87.5	71.2	18.3
DF 261 W	W								
DF 271 W	W								
DF EX 2101 R	R								
DF EX 2102 R	R								
DF EX 2103 R	R								
DF EX 2104 R	R								
Dyna-Gro 9002	R	9.2	8.6	68.0	61.4	7.0	101.0	64.7	18.9
Dyna-Gro 9070	R	9.1	28.2	66.7	57.7	7.4	114.5	67.1	19.2
Dyna-Gro 9082W	W	10.2	45.9	66.6	53.0	8.4	99.2	71.3	17.9
Dyna-Gro 9120	R								
Dyna-Gro 9151	R	9.8	24.0	64.4	58.3	8.1	139.4	76.2	18.2
Dyna-Gro 9172	R	9.9	20.6	68.5	59.7	7.7	110.1	67.2	19.1
Dyna-Gro 9182	R	9.9	32.3	68.0	53.7	8.1	106.5	67.7	18.7
Dyna-Gro 9242W	W	9.5	28.6	67.2	57.7	7.5	102.5	66.8	19.3
Dyna-Gro WX19799W	w								
Dyna-Gro WX20734	R								
Dyna-Gro WX20738	R								
Dyna-Gro WX21791W	W								
Erisman	R	9.3	20.9	68.6	59.9	7.2	93.9	69.2	19.5
Harbor	R	10.7	23.2	69.6	50.0	8.6	103.7	68.1	19.5
Haubert	R								
HS338R	R	9.5	29.2	69.5	57.1	7.4	107.4	69.3	19.4
ISF 1001	W								
ISF 1115	W								
ISF 790	R								
Jupiter	W	9.2	18.7	70.0	59.7	7.0	91.2	68.4	18.9
KWS305	W								
KWS308	W								
KWS316	W	8.9	21.8	68.5	55.5	7.3	102.4	67.3	19.1
KWS317	W								
KWS319	W								
KWS327	W								
KWS340	R								
KWS356	R								
KWS361	R								
KWS375	R								
LCS3334	R	9.0	21.7	68.6	58.9	7.1	96.1	69.6	20.0
	11	3.0		1 00.0	55.5	,. <u>+</u>	30.1	55.0	20.0

Table 5. Milling and baking qualities.

Ţ.		NIR Kernel		Adjusted Flour		Flour		Sodium	Cookie
		Protein	SKCS Kernel	Yield	Softness	Protein	Lactic Acid	Carbonate	Diameter
Line	Color	(at 12%)	Hardness	(%)	Equivalent (%)	(at 14%)	SRC (%)	SRC (%)	(cm)
		` ′				· ·			
LW2068	R	9.1	19.9	70.0	60.1	6.9	90.7	69.5	19.1
LW2148	R								
LW2169	R								
LW2958	R								
MCIA Flipper	R	9.4	31.2	71.2	55.0	7.2	89.7	68.2	18.9
MCIA Jonah	R	9.2	14.5	69.6	62.7	7.0	102.3	67.4	19.3
MCIA MARLIN	R	9.9	28.8	68.8	55.9	7.6	101.5	70.5	19.1
MCIA Red Dragon	R	10.0	19.2	69.1	57.0	7.8	116.4	68.8	18.8
MCIA Whale	R	9.9	29.8	67.3	54.8	7.7	98.4	71.6	19.0
MCIA Wharf	R	9.7	18.9	67.4	54.8	7.5	86.3	66.5	18.8
MI14W0190	W								
MI16R0720	R								
MI16R0898	R								
MI16R0906	R								
MI16W0133	W								
MI16W0528	W								
MI17R0357	R								
MI17W0224	W								
MI17W0235	W								
MI18R1605	R								
MI18W0200	W								
MI18W0286	W								
Moonlight	W	11.1	28.5	68.7	51.3	9.0	108.3	68.0	18.8
Prestyn	R								
REXP1215	R								
RS 902	R	9.9	21.0	69.9	61.2	7.8	102.9	66.5	19.9
RS 912	R	9.8	32.1	68.1	53.5	7.8	105.9	68.4	19.5
RS 961	R	10.2	71.2	70.8	42.1	8.9	91.4	84.6	16.9
RS 977	R	10.3	26.4	68.5	50.5	8.2	92.3	66.5	19.4
RWEXP1212	R								
Sunburst	R	10.0	49.7	64.2	49.1	8.3	98.3	75.4	18.0
SY 100	R	9.1	19.0	68.4	59.3	7.2	98.0	67.7	19.6
SY 547	R	9.6	40.4	66.2	52.2	7.8	94.0	72.3	18.6
SY 576	R	9.9	36.4	64.9	56.5	8.3	98.6	75.5	18.7
SY Viper	R	10.3	23.3	66.0	55.0	8.3	106.7	78.7	18.5
Tyson	R								
W 300	R								
W 304	R	9.3	19.3	70.3	62.8	7.3	93.3	67.2	20.2
W 305	R	10.2	71.0	71.0	40.9	9.1	91.2	83.4	17.3
W 310	R	10.0	12.5	68.3	51.9	7.7	89.8	67.2	19.1
W 313	R	10.4	33.2	67.7	53.4	8.5	108.2	68.6	18.7
W 322	R								
W 324	R								
Whitetail	W								

Commercially Available Varieties entered in the **2021** Michigan State University Wheat Performance Trials

AgriMAXX Wheat Company	9082W	MCIA Whale
AgriMAXX 2050W	9242W	MCIA Wharf
AgriMAXX 2055W	WX19799W	Moonlight
AgriMAXX 498	WX20734	Sunburst
AgriMAXX 502	WX20738	Whitetail
AgriMAXX 503	WX21791W	
AgriMAXX 505		Michigan State University
AgriMAXX 513	Harrington Seeds Inc.	MI14W0190
AgriMAXX 516	HS338R	MI16R0720
AgriMAXX EXP 2002		MI16R0898
	Irrer Seed Farm	MI16R0906
<u>AgriPro</u>	ISF 790	MI16W0133
SY 100	REXP1215	MI16W0528
SY 547	RWEXP1212	MI17R0357
SY 576	ISF 1001	MI17W0224
SY Viper	ISF 1115	MI17W0235
·		MI18R1605
Albert Lea Seeds	KWS Cereals	MI18W0200
Erisman	KWS305	MI18W0286
LCS3334	KWS308	
	KWS316	Rupp Seeds Inc.
DF Seeds Inc.	KWS317	RS 902
AMBASSADOR	KWS319	RS 912
DF 105 R	KWS327	RS 961
DF 112 R	KWS340	RS 977
DF 119 R	KWS356	
DF 121 R	KWS361	Synergy Ag
DF 131 R	KWS375	Haubert
DF 218 W		Prestyn
DF 261 W	Local Seed Company LLC	Tyson
DF 271 W	LW2068	•
DF 141 R	LW2148	Wellman Seeds Inc.
DF EX 2101 R	LW2169	W 300
DF EX 2102 R	LW2958	W 304
DF EX 2103 R		W 305
DF EX 2104 R	Michigan Crop Improvement	W 310
	Association	W 313
Dyna-Gro Seed	AC Mountain	W 322
9002	Harbor	W 324
9070	Jupiter	WEX 106
9120	MCIA MARLIN	
9151	MCIA Flipper	<u>VCIA</u>
9172	MCIA Jonah	13VTK59-55
		10111100 00

MCIA Red Dragon

9182

Organizations Participating in the **2021 Michigan State University Wheat Performance Trials**

AgriMAXX Wheat Company 7167 Highbanks Road Mascoutah, IL 62258 Phone: 855-629-9432

Agripro 1521 N. Convent St. Suite 200 Bourbonnais, IL 60914 Phone: 815-370-3291

Albert Lea Seed 1414 W. Main PO Box 127 Albert Lea, MN 56007 Phone: 800-352-5247

D.F. Seeds, Inc. P.O. Box 159 905 S. Jackson St. Dansville, MI 48819 Phone: 517-623-6161

Dyna-Gro Seed 4648 S Garfield Rd Auburn, MI 48611 Phone: 989-662-0000

Harrington Seeds, Inc. 2586 Bradleyville Road Reese, MI 48757 Phone: 989-868-4750

Irrer Seed Farm 9621 Dexter Trail Fowler, MI 48835 Phone: 517-719-5710

KWS Cereals 4101 Colleen Drive Champaign, IL 61822 Phone: 330-439-3341 Local Seed Company LLC 802 Rozelle St Memphis, TN 38104 Phone: 901-260-6000

Michigan Crop Improvement Association 2905 Jolly Road Okemos, MI 48864 Phone: 517-332-3546

Rupp Seeds, Inc. 17919 Co Rd. B Wauseon, OH 43567 Phone: 419-337-1841

Synergy Ag 6150 N. Co Rd. 33 Tiffin, OH 44883 Phone: 419-355-6708

Virginia Crop Improvement Association (VCIA) 9225 Atlee Branch Lane Mechanicsville, VA 23116 Phone: 804-746-4884

Wellman Seeds, Inc. 23778 Delphos Jennings Road Delphos, OH 45833 Phone: 800-717-7333